



# Oklahoma

## 2004 School Health Profiles Results

### Selected Topics Fact Sheet

| Health Education  | Among States <sup>*</sup><br>Median | Range          | Oklahoma<br>% CI <sup>†</sup> |
|---|-------------------------------------|----------------|-------------------------------|
| Among schools that required health education, percent that required 2 or more health education courses  | 47.6                                | (16.0 – 76.7)  | 22.8 ± 6.9                    |
| Percent of schools that had a health education coordinator  | 96.3                                | (74.2 – 100.0) | 74.2 ± 3.8                    |
| Percent of schools in which the lead health education teacher had professional preparation in health education or in health and physical education combined | 61.7                                | (13.0 – 90.2)  | 40.1 ± 4.9                    |
| HIV, STD, and Pregnancy Prevention  |                                     |                |                               |
| Among schools that required a health education course, percent that taught HIV prevention   | 96.6                                | (82.1 – 99.4)  | 93.9 ± 4.9                    |
| Among schools that required a health education course, percent that taught STD prevention   | 91.9                                | (73.1 – 96.4)  | 81.6 ± 7.9                    |
| Among schools that required a health education course, percent that taught pregnancy prevention   | 84.1                                | (70.1 – 90.5)  | 73.1 ± 9.2                    |
| Among schools that required a health education course, percent that taught abstinence as the most effective method to avoid HIV infection                   | 94.5                                | (77.5 – 98.5)  | 86.4 ± 7.2                    |
| Among schools that required a health education course, percent that taught how to correctly use a condom  | 37.6                                | (8.8 – 60.0)   | 25.0 ± 8.7                    |
| Percent of schools in which the lead health education teacher received staff development during the past 2 years on HIV prevention                          | 42.7                                | (21.8 – 77.4)  | 77.4 ± 3.5                    |
| Percent of schools with a written policy that protects the rights of students or staff with HIV infection or AIDS   | 59.4                                | (44.9 – 84.2)  | 68.7 ± 4.2                    |
| Asthma  |                                     |                |                               |
| Percent of schools that provided a full-time registered nurse, all day every day  | 36.1                                | (0.5 – 98.4)   | 25.9 ± 3.7                    |
| Percent of schools that obtained and used an Asthma Action Plan for all students with asthma  | 62.2                                | (24.8 – 79.0)  | 42.6 ± 4.4                    |
| Percent of schools that educated school staff about asthma  | 53.7                                | (33.2 – 82.0)  | 54.7 ± 4.0                    |
| Percent of schools that educated students with asthma about asthma management   | 52.4                                | (24.4 – 75.9)  | 33.6 ± 4.1                    |
| Unintentional Injuries and Violence   | Among States <sup>*</sup><br>Median | Range          | Oklahoma<br>% CI <sup>†</sup> |
| Percent of schools that maintained a closed campus where students were not allowed to leave school during the school day, including during lunchtime        | 83.0                                | (43.8 – 96.5)  | 73.8 ± 3.7                    |
| Percent of schools that used staff or adult volunteers to monitor school halls during and between classes   | 88.5                                | (72.6 – 94.3)  | 92.8 ± 2.0                    |
| Percent of schools that required students to wear school uniforms   | 4.3                                 | (0.0 – 24.3)   | 3.6 ± 1.5                     |
| Percent of schools that required students to wear identification badges   | 4.6                                 | (0.0 – 37.7)   | 4.6 ± 1.7                     |
| Among schools that required a health education course, percent that taught accident or injury prevention  | 89.7                                | (77.6 – 96.2)  | 90.1 ± 5.7                    |
| Among schools that required a health education course, percent that taught violence prevention  | 87.6                                | (80.0 – 94.0)  | 90.1 ± 5.8                    |



# Oklahoma

## 2004 School Health Profiles Results

### *Selected Topics Fact Sheet*

#### School Health Councils

|  |      |               |                   |
|--|------|---------------|-------------------|
| Percent of schools that had a school health committee or advisory group that developed policies, coordinated activities, or sought student and family involvement in programs that address health issues | 48.7 | (20.5 – 72.7) | <b>20.5 ± 3.0</b> |
|--|------|---------------|-------------------|

\* Among the 27 participating states with weighted principal survey results and the 25 states with weighted teacher survey results.

† 95% confidence interval.

#### Where can I find more information?

More information about the School Health Profiles is available at  
[www.cdc.gov/healthyyouth/profiles](http://www.cdc.gov/healthyyouth/profiles) or call (888) 231-6405.



U.S. Department of Health and Human Services  
Centers for Disease Control and Prevention